UNITED STATES OF AMERICA

7436 O1 JUN-1 A8:34

DEPARTMENT OF HEALTH AND HUMAN SERVICES FOOD AND DRUG ADMINISTRATION

CENTER FOR DEVICES AND RADIOLOGICAL HEALTH

TECHNICAL ELECTRONIC PRODUCT RADIATION SAFETY STANDARDS COMMITTEE

28th Meeting

Thursday,

May 17, 2001

The Committee was called to order at 8:30 a.m., at the Food and Drug Administration, 9200 Corporate Boulevard, Suite 01-02, Rockville, Maryland 20850 by Chairman Larry Rothenberg, presiding.

PANEL MEMBERS PRESENT:

- DR. LARRY ROTHENBERG, Chairperson
- DR. ORHAN SULEIMAN, Executive Secretary
- DR. DAVID N. LAMBERT, Member
- LTC MICHELE LOSCOCCO, Member
- DR. JOHN M. SANDRIK, Member
- DR. ALICE FAHY-ELWOOD, Member
- DR. MAUREEN MURDOCH NELSON, Member
- DR. MARY V. MARX, Member
- DR. JOHN F. CARDELLA, Member

NEAL R. GROSS

COURT REPORTERS AND TRANSCRIBERS 1323 RHODE ISLAND AVE., N.W. WASHINGTON, D.C. 20005-3701

PANEL MEMBERS PRESENT: (cont.)

MR. ROBERT PLEASURE, Member

CPT JERRY A. THOMAS, Member

DR. WILLIAM R. RICE, Member

DR. GREGORY W. LOTZ, Member

DR. QUIRINO BALZANO, Member

DR. STEVE SZEGLIN, Member

MS. KATHLEEN A. KAUFMAN, Member

PUBLIC SPEAKERS:

MS. LILLIAN GILL

DR. W. HOWARD CYR

DR. RONALD KACZMAREK

DR. TOM SHOPE

DR. STANLEY STERN

DR. ROBERT GAGNE

MR. COLLIN FIGUEROA

DR. RUSSELL OWEN

NEAL R. GROSS

C-O-N-T-E-N-T-S

		PAGE
I.	Greeting and Introduction	. 4
II.	Chairman's Opening Remarks	. 6
III.	Update of Informal Issues	. 13
IV.	Presentation by Dr. Tom Shope Concerns Regarding Radiation Doses from Filmless Technologies	. 39
V.	Presentation by Dr. Stanley Stern on Computed Tomography	. 68
VI.	Presentation by Dr. Robert Gagne on Digital X-Ray Imaging	111
VII.	Open Committee Discussion	178
VIII.	Presentation by Mr. Collin Figueroa on Performance Standards for Non-Medical Products	229
IX.	Presentation by Dr. Russell Owen on Cellular Telephones	250
х.	Open Committee Discussion	292
XI.	Closing Remarks	333

P-R-O-C-E-E-D-I-N-G-S

2

1

(8:30 a.m.)

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

SECRETARY SULEIMAN: I would like to get the meeting started. I am Orhan Suleiman, Executive Secretary for the Technical Electronic Product Radiation Safety Standards Committee, and I want to read something quickly here, and then have introduction and some comments by Dr. David Fiegal, who is the Center Director, and get into the agenda early. Actually, why don't you go ahead and welcome people and I will begin right after you.

DR. FIEGAL: I just wanted to come this morning and welcome you to today's meeting. As Orhan likes to remind me, this is the oldest standing committee now that the T-Board has disappeared. And it is one that we value very much.

You know, in today's agenda, I think it really illustrates the need for continued vigilance and attention to the area of radiological safety, and health, and we will hear today about both very new technologies and some of the issues with safety as to that, and we will also hear about problems we may have

thought have largely been solved.

Ad who would have thought that we would be once again coming back to issues around philosophy, and there have been times when there have been problems with that, and where it seems to increase and the difficulties arising from that.

Also today, we will share with you -Lillian Gill, who is the Deputy for Science for the
Center, will share with you some of our thoughts about
the future of the Rad program and CDRH is one that has
changed over the years, and we appreciate your
thoughts about what the challenges of the future will
be.

There have been focuses that have been taken in the past to assure the safety of these products, but there have been many changes in the some 30 years since the Center for Devices and the Rad Health Program with the Public Health Service merged in a way that they are manufactured, and in the way the whole economy works has made this a very challenging environment.

It is a very challenging environment to be

a national regulatory body in a global economy. Globally, the whole trends are to take down borders and to have free trade, and have rapid interchange together. A lot of our premises of structure in regulatory programs would be that we would have strong input from Rads, and we would have a great deal of national manufacturing, and things which have rapidly changed. So let me again welcome you and thank you and advise us, and to help us.

for taking the time from your busy schedules to come It is important not only for your expertise, but also that this be a public process, and it is a process that has a record, and one that has developed over time.

And I told Orhan that even though I had a meeting across town, I did want to stop and thank you for coming and wish you a product day. Thanks very much.

SECRETARY SULEIMAN: Thank you, Dr. Let me read the formal statement that I need In accordance with the Radiation Control for to make.

NEAL R. GROSS COURT REPORTERS AND TRANSCRIBERS 1323 RHODE ISLAND AVE., N.W.

WASHINGTON, D.C. 20005-3701

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

Health and Safety Act of 1968, Public Law 90-602, the Secretary of the Department of Health and Human Services has established a Technical Electronic Product Radiation Safety Standards Committee, known as TEPRSSC, for consultation on matters relating to technical electronic product radiation safety.

As specified by Public Law 90-602, the Committee consists of 15 members, including the Chairman, who are appointed by the Commissioner of Food and Drug, with overlapping terms of four years or less.

Five members are selected from governmental agencies, including State and Federal Governments; five members from the affected industries, and five members from the general public, or at which least one shall be a representative for organized labor.

Members must be technically qualified by training and experience in one or more fields of science or engineering applicable to electronic products, radiation safety, and standards.

The primary function of TEPRSSC is to

NEAL R. GROSS COURT REPORTERS AND TRANSCRIBERS 1323 RHODE ISLAND AVE., N.W. WASHINGTON, D.C. 20005-3701

provide advice and consultation to the Commissioner of Food and Drugs on the technical feasibility and reasonableness of performance standards for electronic products; to control the emission of electronic product radiation from such products; and to review amendments to such standards before being prescribed by the Commissioner.

The committee is not requested to review individual applications for particular products, specific products. Public Law 90-602 and legislative history clearly indicated that the TEPRSSC members are expected to represent a very wide range of interests, with at least one-third of the committee nominated by the regulated industry itself, and appointed on the basis their being able represent industry-wide concerns.

Section 534 of the Federal Food, Drug, and Cosmetic Act specifies that TEPRSSC members are not to be considered officers or employees of the United States for any purpose, including conflict of interest determinations.

However, to be consistent with FDA's

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

general policies regarding advisory committees, the agency believes that a public disclosure memorandum should be made a part of the public record, which identifies each member and provides their appointment affiliation.

Approved on September 15th and September 23rd, 1998, August 30th, 1999, and June 9th, 2000, be delegated authority of the Commissioner of Food and Drugs, the members of TEPRSSC are -- and I will read their names quickly here.

Representing the General Public, John Cardella, M.D., State University of New York, Syracuse Health Science Center; Mary Marx, M.D., University of Southern California; Robert Pleasure, J.D., The Center to Protect Workers' Rights; William Rice, M.D., American Radiology; Lawrence Rothenberg, Memorial Sloane-Kerric Cancer Center.

Representing the Government is Kathleen Kaufman, Los Angeles County Department of Health Services; Michele Loscocco, Lieutenant Commander, Joint Readiness Clinical Advisory Board; Gregory Lotz, Ph.D., National Institute for Occupational

Veterans Administration Medical Center; Captain Jerry 2 Thomas, Uniformed Services of the Health Sciences. 3 And representing Industry are Quirino 4 5 Balzano, Ph.D., Motorola Florida Laboratories, 6 recently retired; Alice Fahy-Elwood, with Lucent 7 Technologies, New Jersey; David Lambert, 8 Lambert Systems; John Sandrik, Ph.D., General Electric 9 Medical Systems; and Steven Szeglin, PGW, a New York Corporation. 10 I welcome you here, and I would like to 11 12 pass off to Dr. Larry Rothenberg, who is the Chairman. 13 CHAIRMAN ROTHENBERG: Thank you, Orhan. I would also like to welcome everyone here. I would 14 15 like to thank the members of the committee for taking 16 time out from their busy schedules to be here, and to the members of the Center staff who have prepared and 17 18 organized the meeting for this morning. 19 I thought it would be a nice idea to start out and if we could just go around the table and have 20 each of the committee members say a second or two 21 22 about their activities and interests, and I will start

Health and Safety; Maureen Murdoch Nelson, M.D.,

out.

I am a medical physicist at Memorial Sloane Kerric Cancer Center, with a primary interest in activities related to diagnostic radiology, and in particular computed tomography and mammography, as well as other areas,

DR. LAMBERT: I am David Lambert, and I am a Professor in Electrical and Computer Engineering at Carnegie-Mellon University, and my primary interests are in the field of magnetism and electromagnetism, and I work extensively in the field of data storage.

DR. SANDRIK: John Sandrik, G.E. Medical Systems, and I am a medical physicist, working primarily in mammography right now, with some responsibility in general diagnostic radiology.

DR. ELWOOD: I am Alice Fahy-Elwood, with Lucent Laboratories, and I am primarily involved with non-ionizing radiation safety associated with wireless and optical products.

DR. NELSON: I am Maureen Murdoch Nelson, and I am affiliated with the Minneapolis VA Medical Center, and my interest is in research design and

NEAL R. GROSS

11	evaluating research.
2	DR. MARX: I am Vicky Marx, a
3	interventional radiologist, at U.S.C., in Los Angeles,
4	and I represent the Occupationally exposed.
5	DR. CARDELLA: Good morning. My name is
6	John Cardella, and I am currently the Chairman of
7	Radiology at the State University of New York, and by
8	training, I am an interventional radiologist, and most
9	recently have been interested in standards writing for
10	radiation safety, and new equipment and design.
11	DR. RICE: Good morning. I am William
12	Rice, a diagnostic radiologist, in Baltimore, American
13	Radiology. I am particularly interested in
14	mammography.
15	DR. LOTZ: I am Greg Lotz, with CDC's
16	National Institute for Occupational Safety and Health,
17	in Cincinnati. I have a background in biophysics and
18	physiology, with a particular interest towards
19	biological effects of non-ionizing radiation.
20	DR. BALZANO: I am Quirino Balzano,
21	formerly with Motorola, and I was involved for a long
22	time in the design of cellular telephones, and my main

interest was with imagery and radio frequency and 1 laboratory. 2 3 DR. SZEGLIN: Steve Szeglin, PTW. I am a medical physicist, and my main area of interest is 4 ionization chamber, electrometer design. 5 6 MS. KAUFMAN: I am Kathleen Kaufman, and 7 I am the Director of the Los Angeles County Health's Radiation Management Program, and we assure compliance 8 9 with regulations and radiation safety for 350 radioactive materials licenses, and about 18,000 x-ray 10 11 tubes. 12 CHAIRMAN ROTHENBERG: Again, I would like 13 to thank you all for being here. Our next item of business is Ms. Lillian Gill, who is the CDRH's deputy 14 director for science, will give us an update of 15 informal issues. 16 17 MS. GILL: Good morning. I have not been 18 formally knighted the deputy director for science. 19 am acting in the place of Dr. Jacobson, who as we heard recently is retiring from the FDA soon. 20 21 I want to give you an update on many of 22 the issues that were under discussion before this

panel not only last year, but some of these were 1 discussed in previous years as well. 2 3 These will be -- I will be giving you sort of a status report on the fluoroscopy amendments, the 4 proposed amendments to the laser standard, the sunlamp 5 issues on the sunlamp standard, people scanners, the 6 7 t.v. receivers, and ultrasound. And I would also like to give you just a 8 9 brief update on where we are with our revitalization project for radiological health issues. Although our 10 11 agenda doesn't call for discussion of these issues, our subject matter expects are available for questions 12 and are certainly here throughout the day. 13 In the area of the fluoroscopy amendments 14 15 to the CDRH performance standard for diagnostic x-ray 16 efforts systems, our to publish the proposed 17 amendments to the performance standards continue. 18 These amendments, primarily addressing fluoroscopic x-19 ray systems, have been discussed as I said before in 20 prior meetings. 21 Since the June 2000 meeting, since last 22 year, our working group developing this proposal has

completed the required draft impact assessment statement which focuses on the potential costs and benefits of the proposed amendments, as well as on the effects οf the proposed action onthe human environment.

In July of last year that draft assessment was provided to the industry and posted on our website with a request for comments. We received no substantial comments of this assessment. So we refined the document and incorporated into the draft notice of proposed rule making some of those refinements.

In December of last year, the Center approved the draft Federal Register notice, which was reviewed by the Office of Chief Counsel in FDA and commented on. We are in the process really of incorporating their comments, after which the document will be returned to them for final clearance.

It will undergo some further editing. We will need the signatures of the Commissioner's Office, and then we will have and hopefully receive final clearance at the department level.

NEAL R. GROSS

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

So although the proposed amendments are not yet published, the publication of the notice of 2 3 proposed rule making is anticipated this year, for which we will have 120 day comment period. 4 After this deadline the comments will be 5 considered by Agency staff, and will proceed with the б final rule, which would become effective one year 7 8 after its publication. 9 onour current time line, 10 proposed rules should become final sometime in early 2003. And on the issue of our proposed amendments to 11 the laser standard, for your last meeting, Mr. Jerry 12 Dennis provided you with a preliminary draft of these 13 14 proposed amendments. 15 The purpose of these amendments, if you 16 will remember, was to harmonize many 17 requirements of the FDA standard with the pending 18 revisions to the International Electrotechnical 19 Commission, or the IEC Standard 60825-1, and with a 60601-2-22 Standard. 20 At that time the amendments to the IEC 21 22 60825-1 were out for a vote. Your advice to us was to

continue without waiting for the IEC ballot. Your reasoning was at that time that even if the IEC amendments were not approved, they would probably be included with similar effect in the next IEC ballot.

As it turned out the IEC amendments were unanimously approved in October of last year and published this past January. If you will remember the focus of those amendments was the incorporation of some recent photobiological scientific data, and the creation of a new hazard classification scheme for laser products.

The cornerstone of that was the restructuring of the hazard classes and really the creation of a new product hazard class that takes into account the design of the optical instruments that are used to view laser radiation.

Since your last meeting the draft proposed CDRH amendments, and the preamble, have been completed, as well as a concept paper. A guidance document has also been drafted for use while our amendments are going through the process of approval.

This guidance states that we will not

NEAL R. GROSS COURT REPORTERS AND TRANSCRIBERS 1323 RHODE ISLAND AVE., N.W. WASHINGTON, D.C. 20005-3701

object to laser products that do not comply with some 1 of our requirements if they comply with the comparable 2 3 requirements of the IEC standards. such as classification. 4 5 In the area of sunlamp standards last 6 year, we presented five possible changes to the FDA performance standard for sunlamp products. 7 presentation came after a review of comments that were 8 9 obtained as a result of our publishing an advanced 10 notice of proposed rule making. A review of that ANPRM and its comments, 11 12 I believe, are included in your briefing package. FDA 13 concluded that some of the possible changes that were 14 presented in that ANPRM needed more research data, and 15 more analysis before formal presentation before the Board, or the advisory panel. 16 I'm sorry. 17 Therefore. we narrowed our 18 changes to five. One, the establishment of the 19 existing recommended exposure schedule as part of the 20 performance standard itself. 21 Two, the use of the human cancer action 22 spectrum in a manner similar to that used by the

and

International Electrotechnical Commission. Three, requiring those that make significant changes that affect performance of sunlamps, or sunbeds, to assume the responsibility of manufacturers. Four, to require a simpler, easier to read, warning label; and the last, to require warning labels in catalogs, specification sheets, manufacturer's brochures. FDA felt that these changes would be the easiest implement and would be uncontroversial.

relatively At our last meeting, at the last TEPRSSC meeting, representatives from the indoor tanning industry, if you will remember, disagreed with our proposals, suggesting that more analysis should be done before proceeding with any specific proposals.

TEPRSSC advised FDA at that meeting to meet with the affected industry to discuss these and any other proposals to better understand the issues before returning to the panel with another round of possible changes to our performance standard for sunlamp products.

Since that time the FDA has met with

NEAL R. GROSS COURT REPORTERS AND TRANSCRIBERS 1323 RHODE ISLAND AVE., N.W. WASHINGTON, D.C. 20005-3701

1

2

3

4

5

6

7

8

9

10

77

12

13

14

15

16

17

18

19

2.0

21

members of the indoor tanning industry. We met with them on September 13th, 2000. Each of the issues of concern were discussed in detail, and a better mutual understanding of the issues was reached. Summaries of the proceedings of September 13th open meeting have been written by the industry trade journals, and are on several industry related internet sites. We also having additional opportunities for the exchange of scientific information during the upcoming months. On June 7th and 8th, we have a meeting of the national and Federal councils on skin cancer prevention at NIH.

Αt that time, medical, academic, government, and non-government representatives will consider research, regulation, and educational efforts pertaining to skin cancer.

Following this, on June 9th and 12th, is the effects of light symposium in Boston, where FDA will present results of its research on methods in measuring changes human skin following UV exposures.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

Then in mid-July, our experts will attend the American Society for Photobiology's 29th annual meeting in Chicago. Dr. Janusz Beer will participate in the session, entitled, "Burning, Tanning, and Typing."

And Mrs. Sharon Miller and Dr. Howard Cyr will address optimizing exposure schedules and sources. Meanwhile, we will continue to evaluate the numerous responses that we received from the ANPRM and from any subsequent meetings with the medical community and the indoor tanning industry.

And then propose specific amendments to the current performance standard. These amendments will be based on the current best science of the documented bio-effects of UV. These amendments will also bring the current regulations up to date.

However, as we all realize, the regulatory process is never static. New amendments will in-turn also be subjected to change and a better understanding of the science of UV effects and better assessment of the risks associated with UV exposures to sunlamps are developed.

1 On people scanners, the FDA doesn't 2 currently have mandatory standards for security screening systems that utilize ionizing radiation. 3 This is a new technology applied to an old concept as 4 the intentional exposure of people for non-medical 5 purposes has been considered unacceptable in the 6 7 latter half of the past century. 8 To address the wide spectrum of opinion on 9 the subject, and the lack of guidance for regulators 10

the subject, and the lack of guidance for regulators in the industry, the American National Standards Institute, or ANSI, N-43 committee appointed a subcommittee to draft a consensus standard for the security screening of people.

Two of our Center staff are on this committee, Frank Saraous serves as the Chair, and Dan Cassidy, in the Office of Compliance, is also a member. That committee, the N43.17 subcommittee, has held three meetings since last year's TEPRSSC meeting, and resulted in as many drafts.

The first two meetings were held in Rockville on August 28th and August 29th, and in mid-November, November 13th and 14th. The last meeting

NEAL R. GROSS

11

12

13

14

15

16

17

18

19

20

21

was held at the Health Physics Society, and they are meeting in Anaheim, California, in February.

While in California, that subcommittee also visited the U.S. Customs and observed the operation of a body scanner at the Los Angeles Airport, as well as toured the Rad scan (phonetic) plant in Hawthorne, where security scanners are manufactured.

Radiation surveys of different models of standards are made at both locations. Prior to the subcommittee meeting a presentation on the status of the standard was given at the health physics society media conference, and that talk was well received by those in attendance, and no objections to any part of the draft standards were voiced.

That Anaheim meeting was meant to be the last meeting before submitting a final draft to the main committee. However, holding that meeting on the West Coast and in conjunction with the HPS conference allowed for participation by some attendees not at the previous two meetings.

As a result of that, some topics were

NEAL R. GROSS

revisited and discussed at length, resulting in some additional changes. The main changes consisted primarily of removal of the explanatory discussions from the body of the standard to an appendix, to a new appendix, and the replacement of required operator limits with the discussion of operator doses and pertinent recommendations.

This last change was done and was necessary to be consistent with the OSHA and the Nuclear Regulatory Commission's regulation for allowable doses to radiation workers.

The main work that remains to be done in this area is to incorporate these changes that have been agreed upon by the subcommittee. It is hoped that this work can be finalized by E-mail and we won't have to have another meeting.

If so, the draft will then be submitted to the main N43 committee for comment and balloting. This is somewhat behind our planned schedule, which was December 2000. However, the planned publication date of June 2002 remains the goal of the subcommittee.

In the area of television receiver, and the television receiver standard, we are planning for some collaborative training. Our center and ORA, which is part of FDA's Winchester Engineering and Analytical Center, together with the consumer electronics association, are planning to sponsor a course to train manufacturers' personnel on the Federal requirements for television products under Chapter 5 of the Food, Drug, Cosmetic Act.

FDA and CEA have developed videotapes for the course, which emphasize compliance with the performance standards for television receivers in Section 1020.10 of Title 21 of the Code of Federal Regulations, as well as the procedures for testing products for compliance.

An ultrasound, as you may recall, last year the FDA presented a pilot to the TEPRSSC committee for comment. We proposed the use of a voluntary international consensus standard, along with the medical device regulatory requirements, in lieu of a mandatory radiation performance standard for ultrasound diathermy products.

NEAL R. GROSS

1.8

As an update of where we are on the pilot, we are now working out the internal processes and policies for that pilot. We hope to have a guidance issued following our mandated need to clear it through good guidance practices by next fall when the next batch of standards will be recognized for use by the medical device industry.

And lastly our efforts on revitalizing radiological health per the radiological health program and CDRH, over the past several years, we have briefed you on the progress of CDRH's reengineering projects, including the revitalization of the Rad health program.

Since the last TEPRSSC meeting, we have held an open public meeting in November to discuss our proposed prioritization model and explore other improvements in information exchange, standards, and product testing.

The results of this meeting support our approach to prioritization. They support our role as an information clearing house, which includes establishing Federal agency alliances, training

NEAL R. GROSS

19

20

21

22

alliances, and holding periodic meetings with stakeholders.

It includes a recommendation for simplifying and harmonizing standards, and providing training and guidance documents to supplement or even substitute for some standards, and it includes a recommendation for improvements in product testing, such as an orphan instrument development program, and workshops with test labs to modify and simplify testing.

The summary of the open public meeting is website posted on our under special topics We have had over the past couple of reengineering. months a strategic planning effort underway in CDRH, and many of our reengineering activities, including the Rad health reengineering activities, are being folded into the Center's strategic plan, so that many of these recommended changes become incorporated in daily activities and our plans. concludes my remarks. Thank you.

CHAIRMAN ROTHENBERG: Thank you very much,

Ms. Gill, for your comprehensive update of informal

NEAL R. GROSS

issues and activities. Maybe at this time, do any of 1 the committee members have any questions or comments 2 3 for Ms. Gill? 4 MS. KAUFMAN: Kathleen Kaufman. I have a 5 comment and a question. I wanted to clarify one thing, because on the people security scanners, you 6 had mentioned something about the workers meeting the 7 Nuclear Regulatory Commission's exposure limits for 8 9 occupational workers. 10 And it actually appears that what they are moving towards or what they are at this point advising 11 is that they meet the records for the members of the 12 13 public, and not for occupational workers. 14 But I had a question on the sun lamp 15 proposals. Unfortunately, I read most of this book before I left, but this was one section that I read on 16 17 the plane. 18 So didn't have my previous year's 19 notebook. And I was wondering if they are down to 20 five proposals, but what I couldn't recall is if there 21 are proposals that have now been eliminated, and if we 22 might briefly here what FDA is no longer considering.

MS. GILL: I am going to have our sun lamp 1 expert revisit any that have been eliminated. 2 DR. CYR: Yes, there were. I am going to 3 have to scratch my brain here and try to remember what 4 5 they were, but there were some issues with which we originally proposed, which upon examination were not 6 resolved scientifically, and we are just are going to 7 have to wait for more data to come in. 8 9 And I am trying to remember what some of 10 those are now. I don't have my notes in front of me from then, and this is about 3 years ago. 11 12 MS. KAUFMAN: Well, let me ask question, because was there some discussion previously 13 14 about requiring specific warnings to be provided to 15 customers, to the users of the equipment? words, right now it is talking about warning labels 16 17 and catalogs. 18 DR. CYR: Right. 19 DR. LAMBERT: And specification sheets, manufacturer's brochures. 20 and But thought 21 originally there had been some discussion about what 22 would be provided to the customers.

DR. CYR: 1 Well, that would be nice, but that is something that is done at the State or local 2 3 The actual regulations of the salons, per se, is not in our domain. We regulate the manufacturers 4 5 of the lamps and the tanning beds. 6 MS. KAUFMAN: There was never ever any 7 requirement that specific language be provided to 8 users? 9 DR. CYR: We have worked with the National Council of Radiation Control Program Directors, and 10 have come up with language as part of that group that 11 12 should be given to, let's say, clients of a tanning 13 And it is basically a rewording of the same warning that is on the tanning beds themselves. 14 So that the clients coming in knows what 15 16 the risks are, and reads that, and then perhaps even 17 signs an informed consent statement. I do recall now one of the most contentious issues, and that was would 18 19 there be a melanoma warning, per se. Would there be 20 a warning of melanoma. 21 And the science behind that changed in the 22 last 3 or 4 years. About three years ago when this

came up, that was really a hot issue. There were a couple of major studies that had come out, and one of them was an epidemiology study out of Sweden linking an increase in melanoma with sunlamp use. And a second one was a study involving fish of all things, and in which they had looked at the induction of melanoma or the production melanoma in this particular model of fish, because there are actually very few animals that actually get melanoma. Humans are one of the few that get it, and

And they were concerned because there was far more melanoma production in the UVA range, longer wavelengths of UVA, than one would expect if you had just looked at let's say the action spectrum for erythema.

this particular little fish is also one of those.

So it seemed to be far more effective in producing melanoma than it was in producing erythema. Now, since that time there have been other studies and the issue or the epi story is not as clear cut as that.

There are plenty of other epi studies that

NEAL R. GROSS

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

2.0

21

don't show this connection with sunlamps. 1 There are 2 some that do, but it is just not a clear epidemiological conclusion on that. One thing is, 3 too, that melanoma appears on parts of the body where 4 the sun doesn't shine. So it is not as clear cut as 5 6 this for basal or spring carcinoma. 7 DR. RICE: Has there been any 8 consideration for а training program the 9 operators? Is that a --10

DR. CYR: Yes. We have worked before with the States in training programs, and most of that training has been done by our Office of Compliance on how to train inspectors for the most part. Training of actual salon operators and those who run the machines themselves is again probably more of a State issue.

And we work again with the Council on Radiation Control Program Directors and trying to get those programs set up. The industry itself has been fairly active in the last year or year-and-a-half in trying to get training programs. And they have even talked about certification of operators and of tanning

11

12

13

14

15

16

17

18

19

20

21

So there has been some movement on training 1 in the last year or year-and-a-half. 2 3 MS. KAUFMAN: So thee is nothing that you recall that was eliminated that was something that --5 DR. CYR: Well, the melanoma warning, per 6 se, no. 7 MS. KAUFMAN: A melanoma warning? 8 DR. CYR: Yes, because there was a strong 9 a melanoma warning, per se, into 10 standard, and based on as I said the uncertain epidemiology studies, et cetera, and also critiques of 11 the fish study, and what does it really mean. 12 13 There is differences in skin thicknesses, and differences in repair enzymes, et cetera, where it 14 is very, very hard to interpret what happens in the 15 fish and what happens in a human. 16 17 Since that time also there has been not an actions spectrum of another model, on a mammalian, but 18 actually on almost a single point, the induction of 19 20 melanoma in a rodent model worked on by Ron Layne in New Mexico; and his data showed that melanoma fell 21 somewhere in between what Dick Setlo had found for the 22

And

that

fish model, and what one would expect from looking at 1 2 an erythema action spectra. 3 And so it was not as bad as the fish model, but then again it was not as good as erythema. 4 So the action spectra story is still up there in the 5 6 air as to what is really going on with UVA and 7 melanoma. 8 DR. CARDELLA: I am curious in terms of the ongoing meetings that are being held in between 9 TEPRSSC meetings about this issue of sunlamps. 10 the specific concern is the representation or the 11 information being obtained balanced as the American 12 Academy of Dermatology continues to be involved in 13 this, or is it strictly meetings with industry people? 14 15 Are there meetings originally raised those concerns? 16 17 DR. CYR: Well, we have talked to the 18 Academy of Dermatology, and asked them to some of these meetings, and their participation has not been 19 20 as great as has been the industry.

NEAL R. GROSS

industry is far more interested in it than now the

dermatology community.

21

22

I quess the

with

those

So it has been a little heavy on the industry side. We can make a more concerted effort to 2 make sure that dermatology people know about these 3 4 meetings, and try to send somebody to them. 5 The best way for me to do that is through 6 this Federal Council, because the dermatology community is at those meetings, and I have tried my 7 darnest to tell them about these meetings and what the 8 issues are, and put it into the minutes, and hope that 9 10 somebody shows up. They do, but again not in the same numbers 11 12 or with the same intensity as the industry 13 participated. 14 DR. NELSON: And along that line, I was 15 wondering if maybe members of this committee might participate in some of these meetings. I don't know 16 if that is a conflict of interest or not. 17 18 DR. CYR: Well, that is a good point, and I guess I could send announcements to you to send out 19 to this mailing list. 20 21 SECRETARY SULEIMAN: Orhan Suleiman. Well, they are open. I mean, I don't see why members 22

couldn't participate. I wanted to clarify something,
Howard. A lot of the requirements that people are
concerned about, they are already part of the existing
standard.

We are really talking about upgrading the warning label and upgrading an exposure schedule that only deals with one skin type. So one of the controversies that occurred at the workshop back in -- I think it was September, was some of the industry people were really ignorant, and it was like we were writing standards for the first time.

We have standards already on the books, and we are just trying to upgrade them with the more current science. And the meetings -- yes, when the meetings are, send them to me, and we will send them out and let members of the committee learn about them.

DR. CYR: All right.

MS. KAUFMAN: Excuse me, but was there anything on -- I am trying to recall some years back, but was there anything on trying to have more consistency in the output of the bulbs, and the information that was supplied to the manufacturers on

NEAL R. GROSS

Was

assuring more uniformity between various bulbs? 2 there something on that? 3 DR. CYR: Very much so. That was not one of the five, but since our presentation last year, the 4 industry itself had meetings with State regulators, 5 6 and found out that issues of lamp compatibility, and had to exchange one lamp for another, is probably one of their top most concerns. And I think you will notice that at one of the meetings that we had that I just said FDA-Industry Workshop on Lamp Compatibility, and this is in the planning stages. It was very much in the planning stages. When we are going to Chicago to this meeting of the American Society for Photobiology, we are going to meet with several people from industry. They are participating in this conference in Chicago, and we are going to meet after one of your sessions and talk about when to have this meeting, and what issues to bring up, and where to have it. And we hope that this comes up pretty

maybe as early as this fall, September or

9

10

11

12

13

14

15

16

17

18

19

20

21

October of this year, to talk about those issues. 1 2 MS. KAUFMAN: I think that was the one 3 that mainly I was concerned about. 4 DR. CYR: Yes, this is a very important 5 concern, and it is being worked on very actively. Sharon Miller in our engineering group is heading that 6 effort. She knows more about lamp compatibility issues than I do, and she is not here right now, but 8 9 she will be the one that will be heading that 10 particular part of that effort here. 11 CHAIRMAN ROTHENBERG: Thank you. I just wanted to clarify. You will try to maybe get us on 12 13 the announcement for those meetings? 14 Definitely. I will try to get DR. CYR: 15 you on the announcement for those meetings. 16 CHAIRMAN ROTHENBERG: Okay. Thank you 17 very much. Thank you again, Ms. Gill, also. The next item is open public hearing. We have no one who 18 19 had requested to speak here but we do have a statement 20 from Tom Quinn, which was available to the committee, 21 and also on the table outside, and that statement will 22 be entered into the record for the meeting.

Since 1 we are somewhat earlier than scheduled, possibly we will postpone the break to a 2 later time, and sort of in the middle of the session, 3 and would it be possible for Dr. Shope to give his 4 5 presentation now? 6 So Dr. Thomas Shope will speak on concerns regarding radiation doses from filmless technologies. 7 DR. SHOPE: Good morning. My purpose for my part of this discussion this morning is really to 9 10 introduce the topic and to lay some groundwork for the 11 following two presentations which will get into a little more technical details. 12 What I hope to do is sort of pose the 13 issue and introduce it so that you will have an 14 opportunity to think about the issue while the 15 discussions are ongoing, and then following both 16 17 presentations have a period of discussion where we can 18 get your comments and opinions. 19 In conjunction with this issue of digital imaging modalities, we really don't have a specific 20 regulatory proposal for the committee to review. 21 22 Rather, we are only at the stage of wanting to discuss

with the committee some of our early thoughts and possible approaches to dealing with this issue, and the concerns that we have about it.

And technology willing here, we will have a slid ein a moment. The projector takes a moment to warm up, I guess. The first thing that I wanted to do was just briefly discuss with the committee the role of the committee, and this is also to some extent with the audience here, as to the fundamental role of the TEPRSSC committee, which is to provide the FDA and the Commissioner with advice on proposed regulatory standards.

And the other thing is to note that our FDA responsibilities go somewhat beyond just regulatory standards, and get into some areas where the charter for the TEPRSSC committee is not really specific, but as a group of experts whose opinions we value, we would like to have an opportunity to discuss some of the other possible approaches that we might use in addition to regulatory approaches to deal with this particular issue that we want to discuss.

(Brief Pause.)

. |

- 15

DR. SHOPE: Ι have highlighted here briefly the charge of the TEPRSSC committee, and just to emphasize that the charge for the committee is to advice and consultation provide on performance standards related to controlling the emission of radiation from electronic products. The next slide. The Act though gives the FDA

additional responsibilities beyond just performance standards. We are charged in the act with the ability to be involved with the research, development, and training, and operational activities related to public health issues related to electronic product radiation; to maintain liaison and receive information on present and future electronic product radiation issues; to study and evaluate the emissions and conditions of exposure to electronic product radiation; and to develop, and test, and evaluate procedures and techniques for minimizing this exposure. And some of these things are not really regulatory in nature.

In addition to these activities, the Secretary, and in this case the Secretary of Health

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

to

make

and Human Services, who was the person being described in the Act, and that authority is delegated to FDA, collects and makes available the results of research and other studies related to electronic product radiation. And also is authorized recommendations relating to these hazards in ways that might be used to control them. Next slide. So that is the general overview of our FDA authorities and responsibilities regarding electronic product radiation. In particular, the discussion for this morning is of particular concern that we have tomography, digital radiography,

relating to digital x-ray imaging modalities. And here I have described them as computed and computer radiography, and specific descriptions of these will

be presented by the next two speakers.

But the principal concern we have is the sort of lack of a fundamental limitation that we see with regular film screen imaging, which is too much exposure results in black films, which is an optic lesson for the person making the exposure not to use

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

that much exposure the next time around.

But these digital modalities don't have this fundamental limitation. Our concern is what will be the long term impact of this on patient dose, and are there some concerns here that the FDA ought to be taking some action with regard to, and what actions might be appropriate for FDA in this issue, both in the regulatory or the non-regulatory area.

What is the magnitude of the problem that we are concerned about? Well, unfortunately we don't have any current comprehensive national nationwide data on patient exposures from all these different modalities.

It would be nice to have and in fact we are in some discussions with some of the other government agencies currently about how we might do a better job of collecting this kind of comprehensive national data. But that is a story that is still unfolding

Our nationwide evaluation of x-ray trends program, the next program that we operate in conjunction with the conference of radiation control

NEAL R. GROSS

COURT REPORTERS AND TRANSCRIBERS 1323 RHODE ISLAND AVE., N.W. WASHINGTON, D.C. 20005-3701

directors, is really the only representative U.S. data on patient exposures, and that is limited primarily to one examination per year, and it cycles through a number of select examinations that gives us some idea of the trends. But it doesn't really address all of the various examinations that are performed, particularly the less frequently done examinations. reports though indicate that patient doses from CT are a significant portion the total medical exposure these days that patients get from medical procedures.

And the other observation that one can make is that CT does are large compared to other typical diagnostic exams, and we will talk in a little more detail about that in a moment.

So there are some fairly good data about CT available, and not so much from the U.S., but from some other countries that probably have experience very similar to what is happening here in the U.S.

Data for digital radiography and computed radiography is kind of a mixed picture at this point as to what the impact of these technologies are on the

NEAL R. GROSS

1

2

3

4

5

6

8

9

10

11

12

13

14

15

16

17

18

19

20

21

typical patient dose or the average patient dose.

Clearly these systems are capable of delivering patient doses similar to conventional film screen systems, keeping in mind that film screen systems have a range of exposure capabilities, which we refer to as the film screen speed indicator, and depending on the detail needed in the imaging procedure, various speeds of film screen systems can be chosen, resulting in various levels of patient exposure. Next.

Looking at this digital radiography and computed radiography issue a little bit, we don't have any large scale national surveys on patient doses from these two modalities as of yet.

In fact, the current 2001 next survey we hope -- and that is being designed with the intention of capturing some information on chest examinations performed with digital modalities. We are actually trying to seed the sample selection here to make sure that we do capture a significant number of digital modalities doing chest examinations to get some idea of what the impact on patient dose is from those types

WASHINGTON, D.C. 20005-3701

of equipment.

There is indications from the literature looking at reports from various institutions that have been published that computed radiography doses are comparable to film screen doses, and typically comparable to film screen systems having speeds of 200 or 400, in that range.

Clearly, CR systems, as Bob Gagne will discuss later, can present higher doses, and I am not sure if there is much opportunity to use lower doses, although that certainly is capable from their dynamic range to deal with lower doses and provide imaging capabilities.

There are some reports either from individual facilities or manufacturers that describe the implementations of the various types of digital radiography systems and these present somewhat of a mixed fixture.

Typically, we see that there appears to be a patient dose reduction possible from digital radiography systems, but there is a lot of variability and a lot of it depends here. It is clear task

NEAL R. GROSS

dependent on what exam is being performed and the type of equipment being used.

And you will see that there is a variety of equipment for this type of digital radiography procedures. There are some concerns that we have about certain implementations of digital radiography that are fairly dose inefficient, and we want to make sure that we keep an eye of those issues.

There are reports in the literature of dose reductions from typical film screen systems when you replace those by a digital radiography system, 25 to 50 percent dose reductions from the same clinical task, giving the clinicians the imaging information they require to do an adequate exam.

But these are not data from basically nationwide kind of studies or region wide kind of studies. They are more institutional experience, particular exam experience in a particular institution.

The other concern that we have is for digital radiography and computed radiography, what will be the long term stability of those levels. If

basically

a facility moves from film screen to one of these 1 digital modalities, and gets it set up according to 2 manufacturer's instructions and it 3 is 4 performing equivalent to a 200 or 400 film screen system, and the manufacturer now goes away, and it is 5 now the responsibility of the facility to keep the 6 7 system operating appropriately, what are the quality assurance measures that are necessary to make 8 that the doses don't drift. 9 10

And the concern that I think we have is drifting upward, as opposed to drifting downward here. Downward would clearly, I think, result in some adverse image quality issues with regard to noise, and would probably be noticed.

But if the exposure drifts upward. probably the noise in the images gets better and the facility or the clinicians using the images won't really notice adverse imaging quality.

is But there the potential increased patient exposure when the wide dynamic range, the ability to deal with an exposure that for a film screen system would have resulted in a black

NEAL R. GROSS

11

12

13

14

15

16

17

18

19

20

21

film, and knowing that you are overexposing the patient. Next slide.

Just to talk a minute about our knowledge

Just to talk a minute about our knowledge about CT, I picked up a couple of surveys basically to here from the United Kingdom that shows the growing use of CT and the experience in the United Kingdom.

This was a rather comprehensive survey that has actually been done twice now in the U.K. In 1989, they looked at their nationwide experience in CT and saw that CT was 2 percent of the radiological examinations.

But in looking at the population dose impact of that, saw that it contributed about 20 percent of the collective effective dose in the U.K. And they have fairly good data in the U.K. for the collective dose from the other radiological exams, the non-CT exams, which in the U.S., we don't have quite that handle on the situation currently.

In 1995, they did the same sort of study again, and this is data from the National Radiologic Protection Board, and CT was 4 percent of all the exams, but it delivered at that point about 40 percent

NEAL R. GROSS

of the collective dose, showing a significant increase in the dose coming from CT. Similar data is available in a not quite a comprehensive format from Germany, and it shows about the same sort of thing. A recent paper here in the U.S. just gives the experience from one facility, the University of New Mexico, in Albuquerque, where they did a look at what was going on in their radiology department, and saw that currently CT was about 11 percent of all the exams done. But looking at their department's experience, they thought that CT was delivering almost 70 percent of the dose to patients from radiologic procedures in their facility.

I do not think this includes any nuclear medicine does in the comparison, but Ι absolutely sure of that. Next slide. Just to talk a minute about what this collective dose business or

I just wanted to remind folks that for a long time we were using the quantity of the effective dose equivalent, which came from an ICRP report, and

> **NEAL R. GROSS COURT REPORTERS AND TRANSCRIBERS** 1323 RHODE ISLAND AVE., N.W. WASHINGTON, D.C. 20005-3701

effective dose means.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

recently we changed the terminology and the way of calculating this by changing some of the factors, and it is now effective dose.

But really this is a method for accounting for the non-uniform irradiations that occur in diagnostic radiology, using weighting factors and the risks that occur to various organs, depending upon the amount of radiation that an individual organ receives, and the risk for cancer induction in that particular organ from the radiation.

So it is really just a scheme for saying what would be the comparable effect if instead of getting this non-uniform exposure, what uniform whole body exposure would produce the same risk. And so that is what this effective dose means.

It is an equivalent comparison to just a uniform whole body dose. Next slide. And just to give you an example. This is from the United Kingdom study back in '89, but in the first column there I show the effective dose equivalent from computed tomography and in the second column it is conventional x-ray imaging.

NEAL R. GROSS
COURT REPORTERS AND TRANSCRIBERS
1323 RHODE ISLAND AVE., N.W.
WASHINGTON, D.C. 20005-3701

And it compares this equivalent dose or effective dose equivalent for CT of the head, for chest exams, compared to a chest film, and you notice that we all I think know that a chest radiograph is a rather low dose exposure.

And when you look at the effect of it on the overall effective dose, a chest radiograph is equivalent to .05 mSv whole body exposure. The abdomen CT exam is 8.8 Msv. For those who have trouble converting the numbers, 8 Msv is .8 rads, or 800 rads, I'm sorry. I am getting it backwards, too.

It is 800 millirads, or .8 rads, for the pelvis; and for barium enemas, it is not a conventional radiographic exposure, but it is a fluoroscopic exposure that depends critically the length of time the procedure goes on, and not just the dose required to expose a given film, and often there are multiple films.

But the point here of that number is to show that the CT exams and some of the fluro exams are comparable in dose, at least in the U.K. experience about 10 or 12 years ago. Next slide.

NEAL R. GROSS

2.0

Just to show some recent numbers from Germany, or at least more recent than the U.K. numbers, a slightly different breakout, and here we are using the newer effective dose numbers which came about, and the weighting factors and the risk factors are different based on the 1990 ICRP report.

And using those numbers, in Germany, we see doses for a chest CT exam equivalent to about 2 rads, and almost 3 rads for the abdomen exam, or 27 Msv.

The average there is about one rad equivalent whole body dose from a CT exam, and the frequency of these exams in the German experience is shown in the right-hand column. So, head exams are still a large proportion, and they are the lower dose type procedures. Next slide.

The concern in CT is sort of tried to be highlighted in this slide. When you look only at the volume of tissue that is actually imaged during a CT scan, the actual radiation impinging on that tissue is on the order of 10 to 50 Msv, or 1 to 5 rads, in a typical CT procedure.

NEAL R. GROSS

There are clearly CT procedures and ways of operating CT systems that give doses in excess of 5 rads to the tissue that is actually imaged. If you then convert that dose to the tissue that is actually imaged to the effective dose, which gives you the risk comparison to a whole body exposure, the CT doses range from 2 to 20 Msv, or comparable to a .2 to 2 rads whole body dose.

And the point of the slide is to note that these does from CT are now not requiring orders of magnitude extrapolation from the data that we have, say, from the incidents of cancer in the atomic bomb survivors in Japan, who were exposed to doses only slightly larger than this, and I think the lowest range of doses that they have seen increases in cancer incidents among the atomic bomb survivors is in the 5 to the 20 rad range.

And so from 2 rads in a CT exam, we are not far from that. We are not talking about orders of magnitude extrapolation. There is some controversy about this data from the atomic bomb survivors, the 5 to 20 rads. There are papers that will argue that

NEAL R. GROSS

maybe they didn't really see cancer effects of those doses, and it was more like a hundred rads.

But I think the message here is that there is not a tremendous, several orders of magnitude, extrapolation required here, and that is something that we need to keep in mind when thinking about the CT doses. Next slide.

So our concern about CT can sort of be summed up in three areas. We are concerned about CT techniques when children or small patients actually -- it could be small adults that were imaged. There was some recent publications in the American Journal of Roentgenology that raised this issue.

I don't think this surprises, and Stan will talk a little bit about this. We also know that with the modern CT systems they are scanning faster, and that allows larger volumes of the patient's tissue to be scanned, or to be imaged.

And there is a concern about the growing use in some circles of CT use for screening either to get your cardiac artery conditions screened or for lung cancer detection that is being proposed, and

discussed, and studied. And so we are a little -- we want to keep our eye on this issue as well. Next slide.

For digital radiography and computed radiography that are the digital modes that are for conventional imaging, we want to know more about what the actual dose experience is, and actual use and implementation of these systems.

What is the long term stability of the dose levels from these, and is there a potential for dose creep as you might describe it, and what actions by FDA would contribute to the optimum use of these imaging modalities, both from making sure the image quality stays appropriate, and that the clinical tasks are able to be done, and that the dose is kept under control. Next slide.

That ends my introductory remarks, sort of posing what the issues are here. The next presentation will be by Dr. Stan Stern, who will discuss CT. He will be followed by Dr. Robert Gagne, who will discuss the digital radiography and computed radiography issues a little bit.

NEAL R. GROSS

1 So whether you want to call a break now or continue, I guess I will leave that to you. 2 think there is a need for questions at this point, but 3 if people have them, I guess I could entertain them. 4 5 MS. KAUFMAN: Could 1 ask a question? DR. SHOPE: Yes. 6 7 MS. KAUFMAN: Kathleen Kaufman. looked at the other presentations, it appears that 8 9 they are not focusing on doses as much as your 10 presentation did. So my question is on dose. What kind of doses are you seeing on the faster, newer CT 11 12 scanners? 13 DR. SHOPE: Well, I think the numbers here 14 reflect that. 15 MS. KAUFMAN: Do you think that they are 16 the same? 17 DR. SHOPE: I think they are similar. There is nothing so inherent in the dose per volume 18 image with spiral scanners. 19 Again, we are just 20 analyzing the data from the 2000 CT survey, which when 21 that data is analyzed, give us a better handle on that. 22

1 Clearly the percentage of facilities that have spiral scanning, and the multi-slice scanning, is 2 growing. It is probably 80 percent of the CT systems 3 4 in the U.S. now are that way. And I think for the German data that a 5 large number of those systems did have the multiple 6 slice or the spiral scanning capability. So we will 7 know shortly when we finish looking at the next data 8 9 what our real experience is. I believe that the next protocol called 10 for using exactly the same measurement technique for 11 those units as it did for a more conventional CT 12 13 scanner. 14 And in my program, we actually measure the dose using protocols on every CT scanner we inspect 15 every time we inspect them. And we were getting very 16 different values, lower values, for those spiral CT 17 18 scanners, and it caused me some concern relative to 19 the measurement protocol. 20 That maybe there is something different in those scanners that our instruments aren't picking up 21 all of the dose, or --22

DR. SHOPE: Well, clearly, doing dose assessment for the spiral scannings requires a little 2 bit of care in how that is done. 3 There is nothing inherent in a spiral scanning mode that implies that 4 the does are going to be dramatically different, 5 except the implication of the pitch. 6 7 You know, how fast does the patient move through it, and if the patients move through it so 8 that the pitch is much greater than one, obviously the 9 doses are going to be reduced. But you are going to 10 11 be probably having noiser images. 12 So I think we are in the process of trying 13 to sort that out with some of the activities that are 14 ongoing with the IEC standard for CT that will be 15 touched on briefly in the following presentations. 16 CHAIRMAN ROTHENBERG: I think I would just comment from my experience in looking at some of these 17 18 machines that in general the doses are not higher with 19 the spiral scanners, and in some cases they are 20 somewhat lower. 21 But in general one of the concerns is what 22 you mentioned, is covering larger volumes of the

patients. So more tissue would be eradicated, but the 1 dose to the eradicated tissue was similar. 2 Also, some of the screening techniques 3 employ lower techniques, and so I think maybe Dr. 4 Stern will be covering some of that. 5 6 DR. SHOPE: The point about even if you are using sort of the same dose per scan, but by 7 scanning more tissue that raises the effective dose, 8 9 because there are more organs being exposed, and that 10 risk gets factored effective into the dose calculation. 11 12 DR. NELSON: I was wondering if you could 13 give us a comparison between the doses of a helical CT for screening for pulmonary embolisms compared to the 14 VQ scans as a real movement towards going to CT scan 15 instead of VQ scans, and if there is a safety 16 difference, that would be very useful to know. 17 18 DR. SHOPE: I don't think we have any real 19 firsthand experience on that. I think typically in 20 the screening modes that people encourage the reduced MA during those scannings. 21 So that of itself would 22 reduce the dose because the lungs less are

attenuating, and you can get a comparable noise image 1 with less MA. 2 3 So if facilities are doing that, the doses from the lung screening kind of procedures would be 4 lower than the usual CT protocols would result. 5 I don't think we have a lot of specific data in hand. 6 We could take a look at that and try to 7 look at that in more detail, but I think the amount of 8 lung screening is small now, but there is a potential 9 for it to grow as the clinical community learns how 10 11 best to use that modality. 12 CHAIRMAN ROTHENBERG: The two protocols 13 that I have seen in institutions in New York have significantly lower MA, and I would estimate typical 14 15 doses on the order of maybe 2/10s of a rad, or 2 Msv. 16 SECRETARY SULEIMAN: I would like to 17 clarify something. Dr. Stan Sterns' presentation is 18 going to go into a little bit more detail. 19 you are absolutely right that there are a lot of changes that have happened with CT. 20 21 I think I can competently say that we are 22 on top of them. I think that some of the questions

are really challenges. We have not figured out how to answer them.

I think some of the innovative things are making patient follow through much easier, and so you are seeing more patients undergoing exams. The dose is probably per helical scanner, I suspect, maybe lower, but there is some other things that the manufacturers have done regarding the equipment.

Again, Stan will probably go into more detail, but that have actually dose implications. So those are going to be covered in the subsequent areas. The other aside, Stan is pretty modest and he doesn't want to say this, but another task that he is involved with this year is he is developing an organ dose handbook that will basically be aimed at the user community.

And which will allow you to calculate the dose from CT and come up with the normalized, homogenized merit of effective dose. So you can compare the doses.

It is not a trivial task, but he is using some of the British data, and some of the next CT data

NEAL R. GROSS COURT REPORTERS AND TRANSCRIBERS 1323 RHODE ISLAND AVE., N.W.

WASHINGTON, D.C. 20005-3701

that we collected last year, and so that is going to 1 be a tool that is going to allow us to answer some of 2 these other questions regarding what are the doses 3 from the pulmonary CT scan, and how does that compare 4 5 with a chest radiograph, and so on. 6 We don't have those answers definitively, 7 and you probably can get different opinions on what 8 the right answer is, but I think we are going to solve that in the next year or two. 9 10 DR. MARX: Dr. Shope, do you have any data 11 in those slides that you have shown with the increased volume Cts, and increased percentage of dose, and how 12 13 many actual people are getting more than one within a 14 relatively short amount of time, or is it just by a CT 15 scan? 16 Because a large -- you know, if you look 17 in any hospital, people have multiple, multiple 18 studies. 19 DR. SHOPE: I would -- we have not looked 20 at that in any great detail. The study from the 21 University of Mexico that New Dr. Mettler 22 colleagues did, did look at that issue of how many

people have multiple CT procedures, and the number of 1 people having second, third, and fourth CT scans all 2 in one episode was not insignificant in my opinion. 3 And so I think there is a concern there. 4 It is not just that a patient gets this one time, but 5 many patients -- and I don't remember the exact 6 7 percentages, but we can find that in the article.

patients they looked at had had four or more CT 9

procedures.

8

10

11

12

13

14

15

16

17

18

19

20

21

22

That is really off the top of my head, and I could be off a little bit on that. But it was a decreasing trend as you went from one procedure to two, to three, to four.

I recall, it was something like five percent of the

So the numbers of patients having that multiple procedures went down. But it was still a significant number still having four more procedures.

DR. MARX: Now, one other point that I just wanted to clarify is that the study that you referred to with the CT pulmonary angiogram actually a very different study than a screening CT

NEAL R. GROSS COURT REPORTERS AND TRANSCRIBERS 1323 RHODE ISLAND AVE., N.W. WASHINGTON, D.C. 20005-3701

looking for lung cancer in completely asymptomatic 1 2 So that is a little bit of apples and 3 oranges. 4 DR. NELSON: And that is --5 DR. SHOPE: As opposed to --DR. MARX: So there you would not be using 6 your low dose study, but you also have to factor in 7 the higher likelihood that you are actually going to 8 make a diagnosis over the --9 10 DR. NELSON: Right. 11 DR. CARDELLA: I just wanted to make a comment just for those that might not be familiar with 12 13 the phenomenon that is occurring, and that is in 14 regard to the use of CT scan as a screening tool. 15 By and large whether you believe that is 16 a good thing or a bad thing, the observation is that is outside of the factors that traditionally 17 regulate the use of examinations. 18 19 Screening Cts, the way that they are 20 marketed in many markets around the United States is that the patient is self-paying for this, and it is as 21 22 though they would go in for it as you would go in to

cancer, and is my abdomen free of cancer, and the 2 payment is from the patient's own pocket. 3 there is not an 4 So insurance 5 oversight of the process, nor is there typically a referring doctor oversight of it. 6 These patients 7 typically will present themselves for screening to in many cases entrepreneurial groups that have set up 8 9 screening programs for the purpose of 10 production basically. So when you talk about screening as Dr. 11 12 Shope was talking about, it is not like screening 13 mammography, where there are some checks and balances. 14 It is an open free for all in many communities. 15 DR. RICE: Ι have point of 16 clarification. If you are looking for pulmonary 17 emboli, a chest CT is a diagnostic tool, and not a 18 screening tool. No more than VQ scans for pulmonary 19 emboli. And in the clinical setting, diagnostic tool, and it is not a screening modality. 20 21 DR. NELSON: My point is that it is 22 unclear which is preferred; the VQ scan, versus a

have your teeth cleaned to see are my lungs free of

helical CT. And if we were to find that the radiation dose for a helical CT is much greater than the VQ 2 scan, that might push clinicians to prefer the VQ scan 3 4 over the helical CT, or vice versa. 5 And if the helical CT has less radiation б exposure, we might start preferring that. Right now it is up in the air which is the preferred procedure. 7 CHAIRMAN ROTHENBERG: Okay. I think maybe 8 we are now approaching our originally scheduled break 9 time. 10 So rather than move on to the next talk, why don't we talk about a 15 minute break, and we will 11 proceed then at about 5 after 10:00. 1.2 Thank you, Dr. 13 Shope. 14 (Whereupon, the meeting was recessed at 15 9:50 a.m., and was resumed at 10:17 a.m.) 16 CHAIRMAN ROTHENBERG: Okay. We are going 17 to start and with regard to the morning schedule, we 18 are going to try and finish up the items on the agenda for the morning, and have lunch at noon, as opposed to 19 20 1:00 p.m., and see if we can move everything forward a bit. 21

Our next speaker is Dr. Stanley Stern, who

22

is going to speak to us about issues related 1 2 computer tomography. 3 DR. STERN: Thank you very much. My talk is going to be in the framework of standardization and 4 5 regulation with regard to the CT asymmetry. have the next slide, please. 6 7 The purpose of this presentation is to brief you and elicit any comments about FDA activities 8 9 and its current thinking concerning recent developments in x-ray computed tomography. 10 11 The themes running through this 12 presentation are grouped into four categories. 13 are radiological practice, rapid technological change, 14 revision of industry standards and development of 15 guidance and regulation. 16 The most recent reminder of the potential 17 impact of CT conduct is a practice that may expose pediatric patients to an excess risk of cancer from a 18 19 larger than needed radiation dose. And I will discuss this situation in 20 2.1 somewhat more detail in the next slide. Another 22 example of radiological practice affecting dose is the

small, but growing, frequency of profusion or contrast 1 phase studies involving a mode often referred to as CT 2 3 fluoroscopy. 4 The common denominator for many of these 5 interventional applications is the repetitive irradiation of one level of the patient's body as the 6

axial position.

Fixed position repeated rotation studies raise the important question of how a new dosimetry standard might quantitatively account for doses incurred in interventional modalities. How does one

x-ray tube rotates many times at a more or less fixed

evaluate dose and in what precisely defined terms.

Examples of rapid technological changing include multi-slice helical scanning and adaptive current modulation. Multi-slice helical CT is here and now. It represents a major advance in single breath hold imaging, which facilitates visualization of small lung nodules.

It enables angiography, fast volume metric scanning, a three dimensional rendering. However, for some scanner models, multi-slice helical CT also

NEAL R. GROSS

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

brings with it a geometrically inefficient use 1 2 I will talk briefly about that issue in radiation. 3 one of the following slides. Adaptive x-ray tube current modulation 4 refers to the capability of a CT unit to reduce or 5 6 increase the x-ray tube current dynamically as the tube rotates around the patient in order to yield the 7 8 least amount of radiation necessary for visualization. 9 It changes the current on the fly to accommodate the patient thickness attenuating the x-10 ray beam, and it offers the prospect of an automated 11 12 way of obtaining an optimal radiation dose. 13 One key take home message is that the 14 repetitively of change in CT practice and technology has left the field so unsettled that not even the 15 nomenclature is standardized. 16 17 This point is particularly important with 18 respect to recent attempts revise to new 19 international safety standard for CT equipment so as to have scanners display an index of radiation dose. 20 21 There are two aspects of standardization 22 that are intimately related to how dose and associated

20

21

22

parameters ought to be defined. The first aspect is a European initiative to have clinical facilities apply what are called reference levels in their quality assurance programs.

In the United States, there is a similar move underway by the American College of Radiology through a test group of the American Association of Physicists in Medicine to introduce what are called reference values into general use by facilities.

The second aspect of standardization is the potential role of the FDA in requiring new performance standards for CT equipment. briefly discuss both aspects. Can I have the next slide, please.

Recently, there has been publicity about CT exams of pediatric patients subject to excessive radiation dose and to the associated risk of premature cancer modality over the course of their lives.

The publicity stems from four related papers in last February's issue of the American Journal of Roentgenology. Their publication comprises a valuable service for public health because

informs the radiology community about a serious appoint 2 problem that it can solve. 3 Here are the two principal messages that 4 these papers overrule. First. when examining pediatric patients, many facilities do not readjust CT 5 scanner parameters after having previously examined 6 7 adults. 8 No readjustment means that smaller. 9 thinner patients probably receive more radiation than is needed for visualization. 10 Second, there is a 11 practical solution. 12 facilities reduce the x-rav tube current according to the patient weight, the dose can 13 14 cut substantially with no loss of clinical 15 efficacy. In this group of papers the perspective by 16 17 Donnelly and Colleagues even provides a technique 18 chart to guide to current reduction. So in the view of CDRH, the problem is one of appropriate use of 19 20 equipment, and the quickest way of dealing with it is to get this message out to the user community. 21 22 The American College Radiology of

published this issue in its February bulletin to radiologists, and that article also mentions their development of a CT facility accreditation program, with a particular component for pediatric patients.

The American College of Radiology accreditation program has been two years in the making, and it is in the final stages of testing, and it will evaluate the ability of a practice to use the minimum amount of radiation needed to produce high quality CT images.

Furthermore, the current American College of Radiology standard for a thoracic CT have specific recommendations for reducing dose to pediatric patients, such as increase the table increment or pitch, or lower the tube current, and use partial scans if appropriate and shorten scan times.

I have emphasized pitch with bold typed face to underscore its importance in the context of standardization of nomenclature. The way that the ACR standard uses the pitch is a good example of the prevalent understanding of how this particular scanner setting may be increased in order to reduce dose, and

NEAL R. GROSS
COURT REPORTERS AND TRANSCRIBERS
1323 RHODE ISLAND AVE., N.W.

WASHINGTON, D.C. 20005-3701

I will talk about that more later.

One significant contribution of CDRH has made in this area has been through the nationwide evaluation of x-ray trends program in collaboration with the Conference of Radiation Control Program Directors.

We have just completed a survey of computed tomography facilities in the United States, and we posed the question of whether dedicated techniques were used for pediatric patients. 43 percent of the facilities responded yes.

The CT survey is based on a nationally representative random sample of facilities, and this particular information about the prevalence of dedicated pediatric technique represents the first quantitative datum on the true magnitude of the problem.

The mortality projection published in the American Journal of Roentgenology was based on an assumption that facilities generally did not use pediatric technique, and now we know that that particular assumption is not true, and that its

modality projection therefore is overestimated.

Nevertheless, it is worth emphasizing that the underlying points of that study are valid and important. Cancer mortality risk associated with radiation dose is higher in children than adults.

Dose can be reduced in pediatric CT, and there is a long way to go in getting all facilities to use appropriate techniques with pediatric patients. The FDA might consider distributing a safety notification about the potential for larger than needed dose in some pediatric CT practice.

Such information would make people aware of the problem, and it would inform them about appropriate CT technique for pediatric patients. We solicit your input on whether to issue a formal notice, and if so, what particular information it might provide.

If the FDA were to take such a step, we would want to be very mindful of conveying a complete context concerning pediatric Cts, and of underscoring its diagnostic efficacy so as to not to dissuade parents from providing their children with examination

1 benefits that generally overwhelmingly individual risk. The next slide, please. 2 3 A central issue in managing radiation dose and computed tomography is that of 4 defining a good representation of dose with respect to 5 6 the radiation emitted by any particular CT system 7 during a procedure. 8 When one thinks of patient dose, the first 9 thought that comes to mind is that of effective dose. Effective dose is a radiological variable frequently 10 used in occupational health and medical physics, and 11 12 it is a whole body dose equivalent of cancer risk associated with radiation detriment. 13 14 However, because effective dose is tissue based, it is practically impossible to measure. 1.5 So 16 for CT, a good representation of patient dose is a quantity that is first easily measurable for any 17 18 particular CT scanner in the absence of a patient. 19 And, second, that nevertheless corresponds 20 reasonably well to the magnitude of energy absorbed by 21 real tissue. And, third, at the same time reflects 22 examination characteristics, such as duration, spatial

extent, and ideally anatomical locations irradiated. 1 2 At. the core of the issue of representation is how technology and practice have 3 progressed beyond the applicability of dose indices 4 and terminology defined 20 years ago with the adoption 5 6 current U.S. mandatory standard for CT 7 equipment performance. 8 Here are the key features of the current 9 standard with respect to those. The most important aspect is that dose information must be 10 provided as documentation by manufacturers to users. 11 12 There is no requirement for a real time display of radiation output. There is no regulatory 13 14 limit to the dose. There is no limitation on how 15 efficiently the x-ray field may overlap the active 16 detector area used to form images. 17 And there is no requirement 18 provision of any so-called automatic exposure control 19 systems. Such a system would be analogous to that used in conventional radiography or fluoroscopy. 20 21 It would optimize the amount of radiation needed for visualization to minimize dose according to 22

the physical girth of a patient attenuating the x-rays sensed by the detectors.

To sum up the only requirement is that manufacturers document typical values of dose. There are no standards requiring physical limitations to the dose or to the extent of the x-ray field.

A dose is characterized by specially defined quantity called the computed tomography dose index, abbreviated CTDI, and because this index figure centrally represents dose and computed tomography, I would like to take a minute to describe CTDI, and to point out some of its features and its limitations. Can I have the next slide, please.

The overarching limitation is that CTDI is defined only for axial scanning. The second figure depicts what happens in a single axial scan. Typically the X-ray source rotates around the patient, and the figure shows a cylindrical phantom, with no axial movement of the patient's support table.

The beam is culminated narrowly in the axial direction, where single image slice can have a fitness T ranging from one to approximately 10

millimeters.

In a tomographic plane depicted on the left as the phantom face, the beam is fan shaped and broad enough to cover the cross-section being imaged. An essential physical aspect of this irradiation geometry is that even though the x-ray field is tightly collimated the axis, once the field begins to penetrate a patient or a phantom, the radiation scatters a great deal, and its axial extent broadens significantly.

The graph on the right shows this broadening in what is called a dose profile. The lower part of the graph represents a single axial scan dose on the ordinate as a function of the axial position on the absicca.

The profile shows how the dose varies. For example, between points A and B, and the central peak corresponds mostly to the primary beam passing through the collimator.

Although the system is collimated to obtain a tomographic section of thickness of 13 millimeters in this example, in fact the dose extends

NEAL R. GROSS

far beyond the 13 millimeter width, to more than a hundred millimeters on either side of the peak.

To represent this phenomenon of distributed dose as a single parameter, CTDI is measured for a single axial scan in an acrylic phantom, and it is defined in terms of an integral of the dose profile over a range that is intended to include most of the scattered radiation contributions, as well the primary radiation.

This range extends over a distance equal to 14 slice widths, from minus 7T to plus 7T. For slices of thickness greater than 7 millimeters, which was typical of axial scanning 20 years ago, the integral includes most of the area under the curve, and is a reasonably good representation of dose.

The parameter "n" in the definition of CTDI is the number of tomographic images obtained in a single scan, and these days "n" typically is 1, 2, or 4. When more than one tomogram is produced in a single rotation, the collimation is widened to accommodate a total scan width corresponding to the product n times T.

1 One can show that CTDI is really particular kind of procedure dose. In axial scanning, 2 usually one does multiple scans covering a clinical 3 region of interest along the z axis. 4 5 If these multiple scans are contiguously spaced by an increment "I" that is equal to the scan 6 7 width Nt, then the dose contributions from the tails of adjacent profiles add up as depicted by the broad 8 dashed curve in the upper part of the graph on the 9

For such a procedure, CTDU is equivalent to the average dose in the central portion of a series of 14 contiguous scans. There can be other kinds of procedures involving multiple axial scans. For example, the increment "I" between scans may be twice the scan width.

In that case the multiple scans are not contiguous. They are spaced apart from one another, and the average dose in the center of a series would be reduced. There can also be procedures where the scans are spaced more closely to one another.

In other words, they overlap. An index

NEAL R. GROSS

10

11

12

13

1.4

15

16

17

18

19

20

21

22

right.

called the multiple scan average dose, abbreviated MSAD, represents the average dose in the central portion of a series of multiple scans, whether the scans are contiguous or not.

And MSAD reflects the dose associated with the scan spacing of the procedure. The equation on the right shows the relationship between the multiple scan average dose and CTDI. When the scan increment "I" equals Nt, this ratio -- well, that ratio, that denominator, is one, and CTDI is equal to MSAD in that case. Next slide, please.

Starting around 1990, CT systems capable of helical scanning were introduced into the market, and since then they have increased in popularity so much that today more than 80 percent of the most frequently used CT scanners in facilities can do helical scanning.

The figure in this slide depicts a helical scanning movement. The patient support table moves along the z axis at the same time that the x-ray tube rotates, and the x-ray field traces out a helical pattern around the surface of a patient or phantom.

NEAL R. GROSS

The message here is that there are no U.S. mandatory performance standards or definitions dedicated specially to helical CT. The U.S. standards in place pertain to axial scanning CT. When manufacturers and users of CT systems measure CTDI or dose profiles, they operate their state-of-the-art helical scanners in an axial scanning mode.

This particular practice is bothersome because one expects that the dose profile for a single rotation in a helical scanning mode to be broader than that for an axial scanning mode.

And with no special Federal requirements to represent helical dose, there may be some dose information missing for the CT user. While the absence of a formal definition of a dose index for helical scanning tends to undermine the legitimacy of CTDI characterizing dose associated with helical systems, there are on the other hand empirical studies showing first that CTDI measured during helical scanning has approximately the same value as CTDI measured during axial scanning.

And, second, that the helical scanning

analog of MSAD can be approximated by CTDI, divided by the pitch. In summary, the accurate representation of a CT dose for helical scanning is ambiguous. For helical scanning, pitch is an important parameter corresponding to how tightly the helix is wound.

With the advent of multi-slice helical scanning, the definition of pitch has become a point of contention between some medical physicists and manufacturers, whose respective interests are dose characterization and management on the one hand, and promotion of multiple image capability on the other.

The traditionally accepted definition of pitch is the ratio of table travel, delta z, per rotation, to the total slice width, Nt, per rotation. And this definition was reflected in the first edition of the voluntary CT safety standard adopted in 1999 by the International Electrotechnical Commission.

According to this definition, when the value of the pitch equals one, the slice widths of the helix are contiguously spaced. For a pitch greater than one the helix is spread out as depicted in this slide, and the dose is reduced.

NEAL R. GROSS
COURT REPORTERS AND TRANSCRIBERS
1323 RHODE ISLAND AVE., N.W.
WASHINGTON, D.C. 20005-3701

And for a pitch less than one the slice widths of the helix overlap one another. So with this definition, pitch can be used as a factor to manage dose, and as indicated by the relationship between helical MSAD and CTDI, one can reduce MSAD by increasing the pitch.

As I mentioned earlier, the current voluntary standard of the American College of Radiology for adult and pediatric computer tomography uses pitch precisely this way as one means to control dose.

Just two years after the first edition, the IEC has recently adopted a second edition of the CT standard containing a new definition of pitch, which may be undergoing further revision at this time.

The new definition refers to only a single tomographic section width T in what may be a multi-tomogram helical scanning mode. This new definition alters an accepted and conceptually accessible representation of the overlap of the image tissue volume, and it may lead to a long period of confusion in the use of pitch for the quantitative estimation of

NEAL R. GROSS COURT REPORTERS AND TRANSCRIBERS 1323 RHODE ISLAND AVE., N.W. WASHINGTON, D.C. 20005-3701

1.3

dose.

Such ambiguity is a prime example of what may ensue in the absence of precise definitions in a U.S. standard. Next slide, please.

Since the introduction of CTDI 20 years ago, there has been a proliferation of indices to characterize different aspects of radiation dose and computed tomography -- CTDI 100, and CTDI 100 central or peripheral, weighted CTDI 100, and normalized CTDI W, dose-length product.

Without getting into the details, I would like to make three points about the parameter proliferation. First, the circumstance that there are so many parameters and that many have ambiguous definitions, reflects how rapidly the technology and clinical practice of computed tomography has changed.

The nomenclature itself has yet to be standardized. The medical physics and radiology communities have not even settled on the terminology, the words that define dose concepts changing to accommodate technological innovation.

NEAL R. GROSS

The second point is that the core of each of these indices is really comprised of a single common parameter designed CTDI 100, and it is highlighted in the red box. It does not show up so clearly on this slide.

CTDI 100 is like CTDI with two significant

cTDI 100 is like CTDI with two significant exceptions. One, the dose profile is integrated over a fixed length, 100 millimeters, used in a commercially available ionization chamber.

And, two, the reference medium for dose is air, the air of the ionization chamber, and not the acrylic material of the phantom that holds the ionization chamber.

CTDI 100 has several advantages over the FDA quantity CTDI as a dose index. First, for tomographic sections of thickness less than 7 millimeters, which is more common for scanning now than it was 20 years ago, the limits of integration bounding CTDI 100 include more of the scattered radiation contributions from the dose profile than does a traditional FDA index CTDI, which is integrated from minus 7T to plus 7T.

NEAL R. GROSS

18

19

20

21

22

There is a relatively larger scattered radiation contribution the thinner the tomographics section, and by including such contributions, CTDI 100 would tend to more accurately represent the dose for thin section configurations than CTDI, which would tend to underestimate the dose.

Second, fixing the limits of integration also makes CTDI 100 much easier to measure with a fixed length ionization chamber than CTDI because the range of integration determining CTDI varies with section thickness T.

Finally, the dosimetric reference medium. air, of CTDI 100 is less energy dependent and more representative of the x-ray energy absorption coefficient in soft tissue than is polymethl methacrylate, PMMA, which is the dosimetric reference medium, as well as the radiation-scattering matrix defining CTDI.

The third major point of this slide is that some of these indices may be better than others as indicators of radiation risk associated with patient dose.

NEAL R. GROSS

For example, CTDI 100 corresponds to a measure of central slice dose at a specified location in the phantom for one particular procedure, namely multiply contiguous axial scans.

CTDI W is a little more general than CTDI 100 because it corresponds to the central slice dose averaged over the entire tomographic cross-sectional area, but still for the same particular kind of multiply contiguous axial scanning.

Dose-length product, DLP, is even more general, because it includes a measure of the entire scanned region that the other parameters lack. A patient receiving a combined chest and abdomen examination incurs a greater risk than one who has a chest exam alone simply because of the larger volume irradiated in the first case.

While CTDI W would have the same value for these two cases, the dose length product would be larger for the combined chest and abdomen exam than for just the chest exam, and therefore the dose length product would better indicate the risk than would CTDI W.

NEAL R. GROSS REPORTERS AND TRANSCRIBE

The bottom line is that when we define dose indices, we should be guided by their intended use. Next slide, please. This slide highlights a dose problem associated with the development of some models of CT systems capable of multi-slice scanning.

The problem has to do with how efficiently

The problem has to do with how efficiently the radiation emitted is actually used to produce clinically useful images, versus how much of it is absorbed by the patient without contributing to the image.

This phenomenon has been characterized as over-beaming by Hans Nagel, and it is a good example of how technological progress may carry challenges as well as rewards.

The figures schematically depict a comparison of the dose profile for a single-slice system versus that for a multi-slice system. In both figures, the length of the active detector along the z axis, the axis of rotation, is shaded green, and both figures portray the same overall width of the tomographic section visualized, 4 millimeters in this example.

1.3

1.8

The difference between them is how that 4 millimeter section is produced. The figure on the left corresponds to a single slice system, and the x-ray detector is a single unit whose active length is wide enough to subtend practically the entire axial distribution of the radiation that has been transmitted through the patient.

This distribution is just like the dose profile that I showed on a previous slide. Here it is represented by shades of gray, as well as by the heights of the rectangles. The central black area, the umbra, for the most part contains the primary radiation that has passed first through the prepatient collimator, and then the patient.

On each side of the umbra are lighter gray areas comprising the penumbra and radiation that has been scattered from its initial direction by passing through the patient.

All of this axially distributed radiation is sensed by the detector, and I am talking about the single slice case. And the detector signal is sent to the computer as part of the process of image

construction.

The full-width at half-maximum intensity, which is denoted FWHM, of 4 millimeters, is determined for the most part by how wide the pre-patient collimator is open.

The figure on the right represents a multi-slice system. In this example, there are four independent x-ray detectors, each corresponding to a width of one millimeter along the z axis.

Depending on how the radiologist wishes to visualize the region of clinical interest, the signals from each of the four detectors may be kept separate to form four separate images, each one millimeter wide, or they may be summed 2 by 2 to yield two separate images, each 2 millimeters wide.

Or all four may be summed to yield one single 4 millimeter wide image. In any case the width overall of the tomographic section or sections visualized is 4 millimeters.

And it is determined not by the prepatient collimator, but by the physical dimensions of the detectors themselves. In order for each detector

NEAL R. GROSS DURT REPORTERS AND TRANSCRIBER

COURT REPORTERS AND TRANSCRIBERS 1323 RHODE ISLAND AVE., N.W. WASHINGTON, D.C. 20005-3701

to separately sense an evenly distributed amount of 1 radiation, 2 some CT models have the pre-patient collimator open wider than 4 millimeters. 3 4 Wide enough to ensure that each detector 5 lies in the umbra of the radiation field, with none in 6 the penumbra. Such a system produces overall an axial distribution of radiation that may be significantly 7 broader than 4 millimeters, 6 millimeters in this 8 9 example. 10 And this penumbra radiation is absorbed by the patient without being used to form the image. 11 all multi-slice CT systems operate this way. 12 But 13 their prevalence is an open question, and so is the necessity of desirability for any system to operate 14 15 with this kind of inefficiency. Is there a sufficient diagnostic advantage 16 17 for such multi-slice systems? Is the patient being 18 exposed to unnecessary radiation? We don't know. Next slide, please. 19 20 just given an overview radiation-dosimetry issues in computed tomography, and 21 now I would like to outline current FDA activities 22

related to CT.

Data acquisition for the CT survey of the Nationwide Evaluation of X-rays trends has just ended, and we are reviewing returns for entry into a database which we expect to complete by September.

Last year before this Committee, I spoke at length about the objectives and design of the CT survey. Preliminary results offer a tantalizing preview of the valuable information that we will gain about CT dose in the United States.

For example, the multi-scan average dose for head exams increased from 46 mGy in 1990 to 54 Mgy in the year 2000, a change of ore than 15 percent. This result and others exemplify how advances in CT scanner technology may contribute to increased collective dose, as well as to improved diagnostic or interventional efficacy.

CDRH has been active in developing consensus standards and guidance at an international level. There are CDRH representatives to the International Electrotechnical Commission maintenance team responsible for the IEC safety standard for CT

NEAL R. GROSS

equipment, to the IEC working group developing an acceptance test standard for CT systems, and to a task group of the American Association of Physicists in Medicine that is developing recommendations for the American College of Radiology about reference values for diagnostic x-ray exams.

The use of reference values or reference levels would constitute a progressive move in an facility quality assurance program of radiation protection. These values correspond to the 75th percentile of the distribution of dose as measured for a particular radiological procedure.

Reference values are norms or benchmarks, to which a facilities practice may be compared. When the reference level is exceeded in any particular examination, the facility may investigate to see if it is possible to reduce exposure without adversely affecting image quality.

So although we critically reviewed the IEC CT standard, and believe that it needs significant clarification and revision, its requirement for display of dose indices represents a major step

NEAL R. GROSS COURT REPORTERS AND TRANSCRIBERS 1323 RHODE ISLAND AVE., N.W. WASHINGTON, D.C. 20005-3701

forward, because displays would be an essential tool 1 2 for evaluation of patient dose. 3 Knowing the dose is the starting point for implementing a reference value program. 4 We are developing as Orhan mentioned, and I guess I am not so 5 modest, but we are developing a Handbook of Patient CT 6 Doses from CT Examinations for medical physicists and 7 8 radiologists. 9 It will be a compendium of doses radiation sensitive tissues for approximately 10 50 11 different CT examinations, and it is designed to be generally applicable for any particular scanner model. 12 13 This handbook is expected to help facility 14 quality assurance programs, and it will facilitate 15 risk communication between clinical staff and 16 patients. 17 There is a research effort underway to 18 develop definitions and quantitative understanding of the relationship between dose profiles, dose indices, 19 20 pitch, and related parameters for helical CT. 21 We expect to report preliminary results of 22 this work at the next annual meeting the

Radiological Society of North America. Finally, the CDRH Office of Compliance continues it oversight with reviews of CT product reports submitted by manufacturers.

For CT, the Office has been focusing recently primarily on issues that may be of pressing concern. For example, the potential problem of overbeaming that I alluded to earlier. Next slide, please.

We believe that many of the concerns related to technological and clinical developments in computed tomographic exposure and dosimetry might be best addressed through a two-tiered regulatory approach; a policy decision that can be implemented in the near term, and mandatory regulations that might go into effect over a longer period.

It should be emphasized that the idea for a regulatory approach is being introduced now for the purpose of public discussion. It is just an idea, and not an announcement of definitive or irrevocable intent, and it represents FDA's current thinking of possibilities only.

NEAL R. GROSS
COURT REPORTERS AND TRANSCRIBERS
1323 RHODE ISLAND AVE., N.W.
WASHINGTON, D.C. 20005-3701

No decisions have been made to implement either a new policy or to propose new rules. 2 3 short term, we are considering the development of guidance for manufacturers that would support the 5 following possible new policy.

> In the information about CT dose provided to users, manufacturers would have the option to specify values of CTDI 100 in lieu of CTDI as it has been previously defined.

> As I described earlier, we believe that CTDI 100 holds advantages of practicability of measurement and fidelity to issue dose. manufacturer were to elect to provide values of CTDI 100, the manufacturer would also need to provide tables enabling a user to convert the CTDI 100 values to the traditionally defined values of CTDI.

> In the long term, regulations might be proposed to resolve the number of the currently outstanding issues that have been discussed in this presentation. The first point is that one would want to update the requirements for provision of dose information.

22

4

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

NEAL R. GROSS COURT REPORTERS AND TRANSCRIBERS 1323 RHODE ISLAND AVE., N.W. WASHINGTON, D.C. 20005-3701

1 Terminology, indices, and ancillary parameters, such a helical pitch, would be clearly 2 defined in terms of conveniently measurable quantities 3 reasonably related to system radiation output apropos a typical procedure done in a given scanning mode. 5 Most likely CTDI 100 would be required for 6 axial scanning, but possibly a newly defined analog 7 might be required for helical scanning. 8 Second, an appropriately defined index or indices of patient 9 examination dose might be required for display. 10 11 The indices displayed ought be 12 sensitive to the particular CT procedure a patient receives, and for that purpose indicates displayed 13 might not necessarily be the same as those required in 14 15 the dose documentation. 16 In other words, we might want dose indices 17 displayed, one, to include the effect of table 18 increment or pitch reducing or increasing dose. to incorporate the range along a patient's body that 19 20 is covered in the scanning. 21 And, three, to reflect the repetitive 22 rotations of a CT fluoroscopic procedure or contrast

study; or we might want to harmonize the displayed 1 indices with the two reference-value parameters 2 required in the European Union, namely dose-length 3 4 product, and CTDI W. 5 In any case, we will face some important 6 practical questions. Who will see the displayed indices of dose? What will they do about what they 7 8 see? 9 the values are displayed at operator's console only, will only the radiological 10 technologist see these values on a regular basis? 11 Will the values be archived in association with exam 12 images? Will the values be reviewed by anyone? 13 14 How can a quality assurance process be facilitated? 15 The third possible regulation is one that 16 might set limits on the axial extent of the radiation 17 field, vis-a-vis the length of the active detector 18 matrix. 19 The issues of over-beaming and efficient use of radiation need to be considered carefully with 20 21 regard optimal to pre-patient collimation 22 diagnostic efficacy.